

A GALAXY X-RAY FUNDAMENTAL PLANE?

M. Fukugita^{1,2} and P. J. E. Peebles^{1,3}

¹*Institute for Advanced Study, Princeton, NJ 08540, USA*

²*Institute for Cosmic Ray Research, University of Tokyo, Tanashi, Tokyo 188, Japan*

³*Joseph Henry Laboratories, Princeton University, Princeton, NJ 08544, USA*

ABSTRACT

We suggest the radii and luminosities of the X-ray emitting halos of elliptical galaxies define a fundamental plane with the star velocity dispersions, as for the corresponding optical observables. Since the X-ray emitting material usually is at larger radius than the stars this can be interpreted as additional evidence for a relation between the space distributions of stellar and dark mass, in analogy to the Tully-Fisher relation and flat rotation curves for spiral galaxies. The picture is complicated, however, by ellipticals with relatively low X-ray luminosities that appear to have modest dark halos, quite unlike the standard picture for spirals.

Subject headings: galaxies: elliptical and lenticular, cD — cosmology: dark matter — X-rays: galaxies

1. The Fundamental Plane

The presence of dark matter in spiral galaxies is well established (under conventional physics) and correlated with the stellar mass, as in the Tully & Fisher (1977) relation between the luminosity in the inner region, which is dominated by stars, and the circular velocity in the outer part that is thought to be dominated by nonbaryonic dark matter. Dark matter in elliptical galaxies is less well explored, because there are fewer accessible probes outside the concentration of starlight, but observations of X-rays, globular clusters and planetary nebulae show an extended dark halo in M87 (Fabricant & Gorenstein 1983) and several other X-ray emitting galaxies (Forman, Jones Tucker 1985; Bridges 1998)

It would be interesting to know whether elliptical galaxies have an analog of the Tully-Fisher relation between the baryonic and dark matter. The fundamental plane (Djorgovski & Davis 1987; Dressler et al. 1987) and the Faber-Jackson (1976) relation apply to properties of spheroids that are likely to be dominated by stars; they seem to have little to say about the dark matter. The X-ray emitting material, that tends to be at larger radii than the stars, has luminosity that correlates with the starlight, roughly as $L_x \sim L_o^{1.8}$, but with an order of magnitude scatter in L_x at given L_o (Fabbiano, Gioia & Trinchieri 1989; Donnelly, Faber & O’Connell 1990; Eskridge, Fabbiano & Kim 1995). Mathews & Brighenti (1998) propose a tighter relation, $L_x/L_o \propto (r_x/r_o)^{0.6 \pm 0.3}$, where r_o and r_x are the optical and X-ray half-luminosity radii. In this paper we present evidence for an even tighter fundamental plane relation in X-ray properties, with a looser connection to optical observables, and we consider possible implications for dark halos.

We write the fundamental plane relation as

$$L \propto \sigma^a r^b, \quad (1)$$

where r is the half-luminosity radius for the optical or X-ray luminosity L , and σ is the central line of sight star velocity dispersion. For our purpose this is simpler than the κ -space convention of Burstein et al. (1997). If the mass-to-light ratio were constant in an isothermal sphere with small core radius the parameters would be $a = 2$ and $b = 1$. For the proposed X-ray relation it would be better to use the X-ray plasma temperature, but that might best await further measurements, perhaps with the CHANDRA satellite.

Scatter plots around equation (1) are shown in Figure 1 for $a = 2$ and $b = 1$, and in Figure 2 for a and b adjusted to minimize the scatter, as discussed below. We choose units for the data set in each panel so $\log L$, $\log \sigma$, and $\log r$ have zero medians. The lines are equation (1). The filled squares are the 11 galaxies for which Mathews & Brighenti (1998) find there are suitable X-ray images from which they can estimate half-luminosity radii. The optical data for this sample are from Bender, Burstein & Faber (1993). The open symbols are the 19 ellipticals with measured X-ray luminosities as well as optical quantities, from the compilations of Eskridge, Fabbiano & Kim (1995) and Bender, Burstein & Faber (1992). The data file we use is available on request.

An optical fundamental plane relation is shown in the upper left panels of the figures. The relation for the X-ray luminosity L_x and half-luminosity radius r_x , with the optical velocity dispersion, is in the lower right-hand panel. The other panels combine the optical half-light radius r_o with L_x , and L_o with r_x .

A least squares measure of the scatter may be unduly affected by non-Gaussian fluctuations, so we use medians. With the normalization to zero median values, a measure of the scatter is the median of

$$A(a, b) = |\log L(i) - a \log \sigma(i) - b \log r(i)|, \quad (2)$$

in the sample of $i_x = 11$ galaxies with X-ray radii or $i_x = 30$ galaxies with X-ray luminosities but not necessarily measured X-ray radii. If L , σ and r were statistically independent the expected frequency distribution of the $A(a, b)$ would be the same as the distribution of

$$B(a, b) = |\log L(j) - a \log \sigma(k) - b \log r(n)|, \quad (3)$$

where j , k , and n are drawn independently at random from the integers $1 \leq i \leq i_x$. The ratio

$$R(a, b) = \text{med } A(a, b) / \text{med } B(a, b) \quad (4)$$

is a measure of the correlation. We use the median of 5000 samples of B , and find the a and b that minimize $R(a, b)$ by numerical search. Table 1 lists the results.

For L_o , r_x there are several minima, none sharply defined; Table 1 and Figure 2 show a representative case. For L_x , r_o there is another local minimum at $a = 0.1$, $b = 1.9$ with $R = 0.43$, almost as small as in the table, but the fit does not look as good as Figure 2. When medians are replaced with means the

minima are broader but at similar positions. With mean square values the minima are still broader and a and b at the minima are substantially different for all but L_o, r_o .

The fifth column in Table 1 is R at $a = 2$ and $b = 1$, for constant M/L . Under the postulate of statistical independence of L , r and σ , the probability that R is less than the data value $R(2, 1)$ is the probability that more than half of i_x random selections of B are less than the median value of $A(2, 1)$. This probability, in the last column of the table,¹ is computed using the binomial distribution and the fraction of the i_x^3 values of B that are less than $\text{med } A(2, 1)$. The probabilities seem consistent with the visual impression of Figure 1: one sees convincing correlation for L_o, r_o , little evidence of it for L_o, r_x , and the impression that the L_x, r_o case could be improved by adjusting a and b , as in Figure 2. The correlation in the L_x, r_x case looks promising albeit not compelling, at about the 3% level. The very discrepant galaxy in this panel, NGC5044, also is the largest anomaly in the radius-luminosity correlation of Mathews & Brighenti (1998). This galaxy fits the optical fundamental plane; the problem is not that the distance somehow is wrong.

Adjustment of the parameters to minimize R for L_o, r_o only slightly reduces the scatter. The scatter is also similar for the optical fundamental plane parameters $a = 1.51$, $b = 0.78$ from Jørgensen, Franx & Kjørgaard (1996). That is, our data set for L_o, r_o shows a reasonably tight fundamental plane relation for a substantial range of values of the power law index of σ , $1 \lesssim a \lesssim 2$, when b is adjusted to a value near unity. The parameters for L_x, r_x are not tightly constrained either, as one sees from the similar quality of fit in the two figures. Our sample is biased by the restriction to galaxies with detected X-ray luminosity, and it may have some distance errors, but the selection does yield a tight relation among the optical observables and a promising relation among X-ray observables.

2. Discussion

A closer test of the proposed X-ray fundamental plane awaits a larger sample, ideally with X-ray tem-

peratures, but the present case seems good enough to motivate preliminary discussion.

The L_o, r_o and L_x, r_x relations in the constant M/L model imply $L_x/L_o \propto r_x/r_o$. This agrees with the Mathews & Brighenti (1998) relation at one standard deviation. We are proposing a relation among the velocity dispersion and the X-ray luminosity and radius that is tighter than similar combinations of σ with L_x, r_o or L_o, r_x .

The optical fundamental plane in our sample is roughly consistent with the picture that the spheroid mass is dominated by old stars with close to universal mass-to-light ratio (as in examples of M/L in ellipticals with cold gaseous disks; Pizzella et al. 1997). Among all high surface density early-type galaxies M/L correlates with M (Burstein et al. 1997 and references therein), but the constant M/L model could be an adequate picture in the mass range of X-ray luminous ellipticals. This model requires close to universal star age and mass functions, to account for near constant M/L dominated by stars, but it says nothing about homology of the spheroids.

From the X-ray data we can argue for the following constraint on homology. The X-ray emitting material, whether accreted from outside a galaxy or in part shed by stars, settles at characteristic radius r_x . The halo circular velocity v_x at r_x determines the temperature T_x . That with the X-ray emitting mass M_x determines the X-ray luminosity L_x . We are suggesting r_x and L_x are tightly related to the star velocity dispersion σ . Since r_x is not tightly correlated with the star half-light radius r_o this requires that v_x is well correlated with σ , meaning that stellar and dark mass distributions are correlated. This is an analog to the Tully-Fisher relation between stellar and dark mass in spirals. Our argument also implies that r_x is tightly correlated with M_x , though both are only loosely related to the optical observables. This is curious enough to be treated with caution, but the Tully-Fisher relation — the “conspiracy” relating baryonic and dark matter in spirals — also is curious. The dark matter “conspiracy” in ellipticals is more directly probed by comparing star velocity dispersions and X-ray temperatures. Davis & White (1996) find evidence for this correlation, and Brighenti & Mathews (1997) for a universal form for the X-ray temperature run. Further exploration with CHANDRA will be of considerable interest.

If the proposed analog to Tully-Fisher is established it is not likely to be the whole story, because

¹A computation of probabilities or of statistical uncertainties of a and b when these parameters are adjusted would be desirable but impractical by the present method because each random pseudo-data set would require a separate parameter search.

some elliptical galaxies seem to be endowed with relatively little dark matter. For M105 Ciardullo, Jacoby & Dejonghe (1993) find $M/L_B \sim 7$ solar units at 10 kpc radius, from the redshift distribution of planetary nebulae. This is consistent with stellar values for an old population without dark matter. Schneider's (1991) analysis of the HI ring that encircles this galaxy and NGC3384 yields $M/L_B \sim 25$ at semi-major axis 100 kpc. Unless this is a near face-on S0 galaxy, and a near face-on ring accidentally matches Schneider's excellent fit of angular position and radial velocity to a Keplerian orbit, this is a marked contrast with what we know about spiral galaxies. Several other examples are in the review by Bridges (1998). In particular, in NGC5128 (the host of Cen A) Peng et al. (as quoted by Bridges 1998) find that if the anisotropy of the planetary velocities is independent of radius the projected mass increases with radius only to 20 kpc, with M/L_B as low as 15 at 50 kpc radius, again not much more than is in the stars. These ellipticals with low M/L are not associated with strong X ray emitting haloes. Well-established examples of elliptical galaxies with prominent dark mass halos, as reviewed by Bridges (1998), tend to be strong X-ray sources.

We have proposed that the dark mass halos of X-ray luminous ellipticals fit an X-ray fundamental plane relation, and that this implies an analog of the Tully-Fisher relation between baryonic and dark matter observables. The evidence is that another class of ellipticals have normal stellar spheroids but low X-ray luminosities and inconspicuous dark matter halos. If confirmed, these are interesting clues and challenges to the theory of galaxy formation.

We are grateful to David Hogg and Penny Sackett for discussions, and to David Burstein and Bill Mathews for their careful reading and helpful comments on our paper. This work was supported in part at the Institute for Advanced Study by the Alfred P. Sloan Foundation and the Raymond and Beverly Sackler Fellowship.

REFERENCES

- Bender, R., Burstein, D. & Faber, S. M. 1992, ApJ, 399, 462
- Bridges, T. 1998, astro-ph/9811136
- Brighenti, & F. Mathews, W. G. 1997, ApJ, 486, L83
- Burstein, D., Bender, R., Faber, S. M. & Nolthenius, R. 1997, AJ, 114, 1365
- Ciardullo, R., Jacoby, G. H. & Dejonghe, H. B. 1993, ApJ, 414, 454
- Davis, D. S. & White, R. E. III 1996, ApJ, 470, L35
- Djorgovski, S. & Davis, M. 1987, ApJ, 313, 59
- Donnelly, R. H., Faber, S. M. & O'Connell, R. M. 1990, ApJ, 354, 52
- Dressler, A. et al. 1987, ApJ, 313, 42
- Eskridge, P. B., Fabbiano, G. & Kim, D.-W. 1995, ApJS, 97, 141
- Fabbiano, G., Gioia, I. M. & Trinchieri, G. 1989, ApJ, 347, 127
- Faber, S. M. & Jackson, R. E. 1976, ApJ, 204, 668
- Fabricant, D. & Gorenstein, P. 1983, ApJ, 267, 535
- Forman, W., Jones, C. & Tucker, W. 1985, ApJ, 293, 102
- Jørgensen, I., Franx, M. & Kjaergaard, P. 1996, MNRAS, 280, 167
- Mathews, W. G. & Brighenti, F. 1998, ApJ, 493 L9
- Pizzella, A. et al. 1997, Astron Ap, 323, 349
- Schneider, S. E. 1991, in Warped Disks and Inclined Rings around Galaxies, ed. S. Casertano, P. D. Sackett, & F. H. Briggs (Cambridge, Cambridge Univ. Press), 25
- Tully, R. B. & Fisher, J. R. 1977, A& A, 54, 661

TABLE 1
PARAMETER FITS

Variables	a	b	R(a,b)	R(2,1)	P(2,1)
L_x, r_x	0.8	0.6	0.14	0.46	0.03
L_o, r_x	2.2	0.2	0.38	1.15	0.7
L_x, r_o	3.9	2.3	0.41	0.79	0.14
L_o, r_o	1.3	1.0	0.18	0.30	0.000018

This 2-column preprint was prepared with the AAS L^AT_EX macros v4.0.




Fig. 1.— Fundamental plane relations with $a = 2$ and $b = 1$ in equation (1). Units in each panel are chosen so $\log L$, $\log r$ and $\log \sigma$ have zero median values.




Fig. 2.— Fundamental plane relations with a and b adjusted to the values in Table 1 that minimize the scatter as measured by equation (4).